

Chapter 20 Worksheet 1 (Ch 20.1-20.5)

Name:


UGA myID:

Instructions:

- Please enter your first and last name as it appears on the eLC roster (do not use a nickname).
- Your UGA myID is a combination of letters and numbers (example: mine is aw00285). **Do not use your 81x number.**
- If you do not have a printer, type your answers in the boxes then upload the worksheet template to Gradescope by **Thursday, January 28th at 11:59 p.m.** Write your work on separate sheets of paper, convert to a PDF and upload to the dropbox on eLC.
- If you have a printer download the worksheet, write your answers and show your work on the worksheet template, convert it to a PDF and upload to Gradescope by **Thursday, January 28th at 11:59 p.m.**

Chapter 20-Part 1

Question 1: For the reaction of the ammonium ion with nitrous acid, the net reaction is




If the initial concentration of nitrous acid is 1.00 M and, after 41.2 s has elapsed, the concentration of nitrous acid has fallen to 0.73 M, what is the average rate of the reaction over this time interval?

- A. 0.0065 M/s
- B. 0.017 M/s
- C. -0.055 M/s
- D. -0.03 M/s
- E. 0.041 M/s

Question 2: For the hypothetical reaction $2\text{A} + \text{B} \rightarrow 2\text{C} + \text{D}$, the initial rate of disappearance of A is $2.0 \times 10^{-2} \text{ mol/L} \cdot \text{s}$. What is the initial rate of disappearance of B?

- A. $8.0 \times 10^{-2} \text{ mol/L} \cdot \text{s}$
- B. $1.0 \times 10^{-2} \text{ mol/L} \cdot \text{s}$
- C. $1.4 \times 10^{-1} \text{ mol/L} \cdot \text{s}$
- D. $4.0 \times 10^{-4} \text{ mol/L} \cdot \text{s}$
- E. $1.4 \times 10^{-2} \text{ mol/L} \cdot \text{s}$

Question 3: Consider the general reaction:



For this reaction, the rate when expressed as $\Delta[\text{Br}_2]/\Delta t$ is the same as

- A. $-\Delta[\text{H}_2\text{O}]/\Delta t$
- B. $3 \Delta[\text{BrO}_3^-]/\Delta t$
- C. $-5 \Delta[\text{Br}^-]/\Delta t$
- D. $-0.6 \Delta[\text{Br}^-]/\Delta t$
- E. None of the above

Question 4: A certain reaction is zero order in reactant A and second order in reactant B. If the concentrations of both reactants are doubled, what happens to the reaction rate?

- A. The rate of reaction is quadrupled.
- B. The rate of reaction is doubled.
- C. The rate of reaction remains the same.
- D. The rate of reaction is halved.
- E. The rate of reaction is quartered.
- D. increase the rate by an order of 9.


Question 5: Consider the reaction $\text{C}_4\text{H}_9\text{Br} + \text{OH}^- \rightarrow \text{C}_4\text{H}_9\text{OH} + \text{Br}^-$. When the concentration of $\text{C}_4\text{H}_9\text{Br}$ is doubled, the rate of the reaction increases by a factor of two. When the concentrations of all reactants and products are doubled, the rate also doubles. What is the overall order of the reaction?

- A. Zero order
- B. First order
- C. Second order
- D. Third order
- E. Fourth order
- F. Fifth order

Question 6: The rate expression for a reaction is shown to be $\text{rate} = k[\text{A}]^2[\text{B}_2]$. If, during a reaction, the concentration of A was suddenly halved and the concentration of B was suddenly doubled, the rate of reaction would


- A. increase by a factor of 8.
- B. increase by a factor of 4.
- C. increase by a factor of 2.
- D. remain the same.
- E. decrease by a factor of 2.
- F. decrease by a factor of 4.
- G. decrease by a factor of 8.

Question 7: Over the time interval 300 to 400 seconds, the rate of reaction with respect to A is $\Delta[\text{A}]/\Delta t = 3.7 \times 10^{-5} \text{ M/s}$. Over the same time interval what is the rate of reaction with respect to B, $\Delta[\text{B}]/\Delta t$?



- A. $\Delta[\text{B}]/\Delta t = \Delta[\text{A}]/\Delta t = 3.7 \times 10^{-5} \text{ M/s}$
- B. $\Delta[\text{B}]/\Delta t = (-1/4)(\Delta[\text{A}]/\Delta t) = (-1/4)(3.7 \times 10^{-5} \text{ M/s}) = 9.2 \times 10^{-6} \text{ M/s}$
- C. $\Delta[\text{B}]/\Delta t = (1/2)(\Delta[\text{A}]/\Delta t) = (1/2)(3.7 \times 10^{-5} \text{ M/s}) = 1.8 \times 10^{-5} \text{ M/s}$
- D. $\Delta[\text{B}]/\Delta t = -(1/2)(\Delta[\text{A}]/\Delta t) = -(1/2)(3.7 \times 10^{-5} \text{ M/s}) = -1.8 \times 10^{-5} \text{ M/s}$

Question 8: For the reaction:




the following initial rate data were collected at constant temperature. Determine the correct rate law for this reaction. All units are arbitrary.

Trial	[A]	[B]	[C]	Rate
1	0.225	0.150	0.350	0.0217
2	0.320	0.150	0.350	0.0439
3	0.225	0.250	0.350	0.0362
4	0.225	0.150	0.600	0.01270

- A. $\text{Rate} = k[\text{A}][\text{B}][\text{C}]$
- B. $\text{Rate} = k[\text{A}]^2[\text{B}][\text{C}]$
- C. $\text{Rate} = k[\text{A}][\text{B}]^2[\text{C}]^{-1}$
- D. $\text{Rate} = k[\text{A}]^2[\text{B}][\text{C}]^{-1}$
- E. None of these

Question 9: The reaction that occurs in a Breathalyzer, a device used to determine the alcohol level in a person's bloodstream, is given below. If the rate of appearance of $\text{Cr}_2(\text{SO}_4)_3$ is 1.24 mol/min at a particular moment, what is the rate of disappearance of $\text{C}_2\text{H}_6\text{O}$ at that moment?



- A. 3.72 mol/min
- B. 0.413 mol/min
- C. 0.826 mol/min
- D. 1.86 mol/min
- E. None of these.

Question 10: The reaction $\text{A} \rightarrow \text{B}$ is first order in $[\text{A}]$. Consider the following data.

Time (s)	0.0	5.0	10.0	15.0	20.0
$[\text{A}]$ (M)	0.20	0.14	0.10	0.071	0.040

What is the rate constant (s^{-1}) for this reaction?

- A. 3.0×10^{-2}

B. 14

C. 0.46

D. 4.0×10^2

E. None of these.

Question 11: A particular first order reaction was monitored over a period of time. The figure to the right summarizes the experimental results where each sphere represents 3.0 mol L^{-1} . The average reaction rate is,


- A. $5.0 \text{ mol/L}\cdot\text{s}$

B. $1.0 \text{ mol/L}\cdot\text{s}$

C. 1.0 s^{-1}

D. $0.20 \text{ mol/L}\cdot\text{s}$

E. 0.20 s^{-1}



Question 12: The decomposition of ammonia on a metal surface to form nitrogen gas and hydrogen gas is a zero-order reaction. At 873°C , the value of the rate constant is $1.5 \times 10^{-3} \text{ mol/L}\cdot\text{sec}$. Determine the number of seconds needed to decompose 90.0% of the ammonia in a solution containing 1.00 grams of ammonia in a 250 mL flask.


seconds

Cuestión 13: Data for the reaction $A + B \rightarrow C$ are given below. Find the rate constant for this system.

Experiment	$[A]_0, M$	$[B]_0, M$	Initial rate, M/s
1	0.030	0.060	2.5×10^{-5}
2	0.030	0.020	2.5×10^{-5}
3	0.060	0.060	10.0×10^{-5}


- A. $2.8 \times 10^{-2} M^{-1} s^{-1}$
 B. $2.8 \times 10^{-2} M s^{-1}$
 C. $2.8 \times 10^{-2} M^2 s^{-1}$
 D. $1.7 \times 10^{-3} M^{-1} s^{-1}$
 E. $1.7 \times 10^{-3} M s^{-1}$

Question 14: At elevated temperatures, methylisonitrile (CH_3NC) isomerizes to acetonitrile (CH_3CN):



The reaction is first order in methylisonitrile. The attached graph shows data for the reaction obtained at 198.9 °C. What is the rate constant (s^{-1}) for the reaction?

- A. -1.9×10^4
 B. +6.2
 C. $+1.9 \times 10^4$
 D. -5.2×10^{-5}
 E. $+5.2 \times 10^{-5}$




Question 15: The following reaction is first order in A and first order in B:



What is the initial rate of this reaction in vessel (b) relative to the initial rate of this reaction in vessel (a)? Each vessel has the same volume. Shaded spheres represent A molecules, and unshaded spheres represent B molecules present at the beginning of the reaction.

- A. rate in vessel (b)/rate in vessel (a) = 1:2
 B. rate in vessel (b)/rate in vessel (a) = 1:1
 C. rate in vessel (b)/rate in vessel (a) = 2:1
 D. rate in vessel (b)/rate in vessel (a) = 4:1



Question 16: The decomposition of a certain insecticide in water at 12 °C follows first-order kinetics with a rate constant of 1.45 yr^{-1} . A quantity of this insecticide is washed into a lake on June 1, leading to a concentration of $5.0 \times 10^{-7} \text{ g/cm}^3$. Assume the temperature of the lake is constant.

$$\ln \frac{[A]_0}{[A]_t} = kt$$

Part A: What is the concentration of insecticide on June 1 of the following year?

g/cm^3 .

Part B: How long will it take for the insecticide concentration to decrease to $3.0 \times 10^{-7} \text{ g/cm}^3$?

years.

Question 17: SO_2Cl_2 decomposes by first order kinetics and $k = 2.81 \times 10^{-3} \text{ min}^{-1}$ at a given temperature. The initial concentration of $\text{SO}_2\text{Cl}_2 = 0.015 \text{ M}$. Determine the half-life of the reaction.

- A. $t_{1/2} = 0.6931 / 2.81 \times 10^{-3} \text{ min}^{-1} = 246.6 \text{ min}$
B. $t_{1/2} = 0.6931 / 2.81 \times 10^{-3} \text{ min}^{-1} = 247 \text{ min}$
C. $t_{1/2} = 1 / (2.81 \times 10^{-3} \text{ min}^{-1} (0.015)) = 2.37 \times 10^4 \text{ min}$
D. $t_{1/2} = 1 / (2.81 \times 10^{-3} \text{ min}^{-1} (0.015)) = 2.4 \times 10^4 \text{ min}$

Question 18: The first-order reaction, $\text{SO}_2\text{Cl}_2 \rightarrow \text{SO}_2 + \text{Cl}_2$, has a half-life of 8.75 hours at 593 K. How long will it take for the concentration of SO_2Cl_2 to fall to 14.5% of its initial value?

hour